



# *Position Statement*

## **Vascular Access for Hemodialysis**

Patients who have reached stage 5 of chronic kidney disease (CKD) or kidney failure with replacement therapy (KFRT) require dialysis or transplant for survival. Patients who require hemodialysis therapy must have a permanent means of access to their bloodstream to achieve this life-sustaining therapy.

### **It is the position of the American Nephrology Nurses Association (ANNA) that:**

- CKD education related to creation and maintenance of the hemodialysis vascular access should be made available to patients with CKD and their families beginning in stage 4. Ideally, as soon as the diagnosis of impending progression into stage 5 is made and need for kidney replacement therapy has been determined, permanent vascular access should be placed.
- Vein preservation of both peripheral and central vessels should be incorporated into patient teaching and care. Blood draws and IV placement should be from/in the dorsum of the hands whenever possible, regardless of arm dominance. Forearm and upper-arm veins suitable for future vascular access should not be used for venipuncture or for placement of intravenous catheters. Subclavian catheters or peripherally inserted central catheter (PICC) lines should be avoided due to the risks of central vein stenosis and occlusion.
- Optimally, all patients requiring maintenance hemodialysis therapy should have a functioning permanent vascular access in place prior to the initiation of hemodialysis. ANNA endorses the recommendations of the National Kidney Foundation (NKF) Kidney Disease Outcomes Quality Initiative (KDOQI) Clinical Practice Guideline for Vascular Access, which recommends a more patient-focused approach with the End-Stage Kidney Disease (ESKD) Life-Plan. The use of hemodialysis catheters greater than 90 days should be avoided if possible to support recommendations by the Centers for Medicare & Medicaid Services (CMS).
- ANNA endorses the recommendations of the NKF KDOQI Clinical Practice Guideline for Vascular Access regarding assessment, access placement, and maintenance.
- A routine Access Surveillance Program should be employed in each dialysis facility to identify and intervene for vascular access dysfunction, enhance long-term access function, and reduce the costs associated with maintenance of access patency.
- Access surveillance and management is an interdisciplinary team function. The patient, nephrologist, nephrology nurse, technician, interventional radiologist/nephrologist, surgeon, and primary care physician should be participants of the team. The social worker and dietitian also can participate in this sustained effort.
- The interdisciplinary team should assure each patient and family has factual information related to all types of vascular access and respect the patient's right to self-determination in choice of vascular access.
- Staff and patient education should include information on infection control measures applicable to hemodialysis access sites.

**Vascular Access for Hemodialysis**

- Staff education should include principles and hands-on cannulation training for vascular access to assure optimal care of the patient's access. Staff education programs should include satisfactory demonstration of knowledge and skills prior to the staff member being allowed to independently perform cannulation. Facilities should adopt best practice models to develop vascular access management plans.
- Education in self-cannulation should be offered to patients (regardless of treatment location) with the ability and access placement location to enable them to take on this responsibility. (This includes in-center, hospitalized, and self-care at-home hemodialysis patients).
- Cannulation of vascular access should be considered as an integral part of successful hemodialysis and an important factor in access outcomes.

**Background and Rationale**

Vascular access for hemodialysis has long been a formidable challenge to those involved in caring for patients with ESKD. In the early 1960s, vascular access for chronic hemodialysis was established using an external device known as the Scribner shunt. This access device consisted of Teflon® tubes surgically implanted into the patient's artery and vein and connected externally by silastic tubing. This method of vascular access was plagued by infection, clotting, and the potential risk of exsanguination should the device become disconnected or dislodged. In 1966, the internal AV fistula was developed by Drs. Brescia and Cimino. The native AV fistula was a significant breakthrough at the time and is still considered to be the "gold standard" for hemodialysis vascular access. In the 1970s, biologic and synthetic grafts became available. In the 1980s, central venous catheters for hemodialysis were developed. Grafts and catheters are associated with a higher incidence of infection and thrombosis; therefore, the native AV fistula is the preferred means of hemodialysis vascular access.

In 1997, the NKF published the Dialysis Outcome Quality Initiative (DOQI) Clinical Practice Guidelines, including Vascular Access Guidelines. These guidelines, revised and expanded in 2000 to include earlier stages of chronic kidney disease, endorse the use of the native AV fistula as the preferred vascular access for hemodialysis. The NKF renamed the project Kidney Disease Outcome Quality Initiative (KDOQI) and the Vascular Access Guidelines were updated in 2006. These guidelines stipulate that when the creation of a native AV fistula is not possible, a graft or central venous catheter may be used for hemodialysis vascular access. These guidelines also stipulate that peritoneal dialysis may serve as a bridge to AVF maturation to avoid hemodialysis catheter placement.

In 2019, the KDOQI Clinical Practice Guideline for Vascular Access adopted a new approach that focuses on the patient's ESKD Life-Plan. The concept of the Life-Plan is to consider a more patient-focused approach to identify modality and access needs from diagnosis to end of life.

Previous collaboration between CMS and the ESRD networks started in 2003 for the purpose of advancing the KDOQI vascular access guidelines. The recommendation from this group increased the goal for AV fistulas to 50% for incident patients and 40% for prevalent patients. In 2005, CMS recognized the work of this group as having a significant impact on the management of vascular access in patients with CKD. The group was renamed the Fistula First Breakthrough Initiative (FFBI) and the AV fistula goal in prevalent patients was increased to 66%. In 2013, this initiative was renamed the Fistula First Catheter Last (FFCL) Workgroup Coalition. The contract between FFCL and CMS has ended.

## **Vascular Access for Hemodialysis**

Nephrology nurses have the primary responsibility to assure the highest quality cannulation to preserve access integrity and prevent access complications. This responsibility includes incorporation of best practices for access cannulation by promotion of expert cannulators and formal cannulation protocols and offering patients the opportunity to self-cannulate. Tools developed by the Making Dialysis Safer for Patients Coalition (which includes organizations such as the Centers for Disease Control & Prevention, ANNA, the American Society of Nephrology, etc.) to improve the safety of vascular access care also should be included.

The following steps may lead to a reduction in morbidity and mortality in patients with CKD: increasing the use of native AV fistulas as the primary option for vascular access; access surveillance efforts to identify access dysfunction; initiation of timely, appropriate interventional procedures; and preserving central and peripheral vessels to assure future access. Implementation of these steps will not only improve the quality of life enjoyed by patients but will also serve to reduce the ever-increasing cost of providing hemodialysis care to patients with CKD.

### **Glossary**

**Access Surveillance Program** – A program developed and implemented by an interdisciplinary group in the dialysis setting to assure that access dysfunction is detected early, so appropriate and timely interventions can be executed. This program should include a combination of routine (at least weekly) physical examinations of the access as well as some form of objective monitoring such as static venous pressures or flow measurements.

**Arteriovenous (AV) Graft** – A tubular biologic or synthetic material surgically implanted to serve as a conduit connecting an artery to a vein.

**Cannulation** – The practice of placing two large bore needles into the AV fistula outflow vein or the AV graft to achieve the necessary blood flow to, through, and from the dialyzer for the duration of the hemodialysis treatment.

**Central Vein Stenosis** – Narrowing of the internal diameter of a central vein, typically associated with trauma to the vessel which results in the development of neo-intimal hyperplasia.

**CKD Stage 4** – Patients with a glomerular filtration rate (GFR) less than 30mL/min/1.73m<sup>2</sup>

**CKD Stage 5** - Patients with a glomerular filtration rate (GFR) less than 15mL/min/1.73m<sup>2</sup>

**Maintenance Hemodialysis** – Routine repetitive hemodialysis therapy required to sustain life.

**Native Arteriovenous (AV) Fistula** – A direct connection between an artery and a vein created surgically to allow higher blood flow through these vessels to facilitate vascular access for hemodialysis. Typical sites for creation of a native AV fistula are the wrist (connecting the radial artery to the cephalic vein) or the elbow (connecting the brachial artery to the cephalic vein).

### **Suggested Readings**

Centers for Disease Control and Prevention (CDC). (2021). Making Dialysis Safer for Patients Coalition. <https://www.cdc.gov/dialysis/coalition/resource.html>

Centers for Medicare & Medicaid Services (CMS), HHS. Medicare and Medicaid Programs; Conditions for Coverage for End-Stage Renal Disease Facilities *Federal Register* / Vol. 73, No. 73 Rules and Regulations; V582. 2008.

**Vascular Access for Hemodialysis**

Counts, C. (2020). *Core curriculum for nephrology nursing* (7<sup>th</sup> ed.). American Nephrology Nurses Association.

Gomez, N. (2017). *Nephrology nursing scope and standards of practice* (8<sup>th</sup> ed.). American Nephrology Nurses Association.

Levey, A.S., Eckardt, K., Dorman, N.M., Christiansen, S.L., Hoorn, E.J., Ingelfinger, J.R., Winkelmayr, W.C. (2020). Nomenclature for kidney function and disease: Report of a Kidney Disease: Improving Global Outcomes (KDIGO) Consensus Conference. *Kidney International*, 97(6),1117-1129. <https://doi.org/10.1016/j.kint.2020.02.010>

Lok, C.E., Huber, T.S., Lee, T., Shenoy, S., Yevzlin, A.S., Abreo, K., Valentini, R.P. (2020). KDOQI Vascular Access Guideline Work Group. KDOQI clinical practice guideline for vascular access; 2019 update. *American Journal of Kidney Disease*, 75(4) (suppl 2):S1-S164. <https://doi.org/10.1053/j.ajkd.2019.12.001>

Medical Education Institute, Inc, (2018). *Core curriculum for the dialysis technician* (6<sup>th</sup> ed).

***Adopted by the ANNA Board of Directors in February 2003  
Revised and/or reaffirmed biennially: 2005-2017  
Most recently revised: August 2021***

***ANNA Position Statements are reviewed and reaffirmed biennially.***