



Position Statement

Nephrology Nurse's Role in Kidney Supportive Care, including Palliative and End-of Life Care

The American Nephrology Nurses Association (ANNA) recognizes that kidney supportive care includes palliative and end-of-life (EOL) care for persons with chronic kidney disease (CKD). This position statement provides nephrology nurses (and other nurses caring for persons with CKD) information, and guidelines related to primary palliative care skills including shared decision-making, treatment preferences, symptom management, advance care planning and EOL care, with a focus on patient quality of life.

It is the position of ANNA that nephrology nurses (APRN and RN) should:

- Contribute to the body of knowledge and direct evidenced-based interventions related to kidney supportive care in individuals with CKD or unresolved acute kidney injury (AKI).
- Engage in conversations regarding kidney supportive care in an environment that promotes informed decision making and respects the individual's goals, preferences, wishes, and values.
- Be involved in policy and advocacy efforts that improve access to equitable kidney care and address the physical, emotional and financial burdens associated with kidney disease and replacement therapies (Diamond, et al., 2021; Gelfand, Mandel et al., 2020).
- Implement an evidence-based approach to symptom assessment and management in all kidney care settings. The use of evidence-based validated tools such as the Edmonton Symptom Assessment Scale-Renal (ESAS-R) or the Integrated Palliative Outcome Scale-Renal (IPOS-R) is important to optimize symptom assessment and management and patients' quality of life (Davison et al., 2024).
- Encourage and participate in the provision of comprehensive kidney supportive care which is holistic and person/family-centered to address social, cultural, psychological, spiritual, religious, and physical concerns.
- Utilize shared decision-making as a fundamental component of patient-centered care, when partnering with individuals and their care partners to determine their preferred treatment options that align with their goals and preferences.
- Develop and maintain competence in providing person-centered kidney supportive care through effective communication, active listening, collaboration with interprofessional teams, utilization of community resources and referrals as a component of their professional and ethical responsibilities in patient care.
- Recognize their personal comfort levels, biases and beliefs regarding supportive, palliative, and EOL care and not allow these to influence a person's decision-making process.

- Be committed to participating in professional discussions, educational programming, evidence-based clinical projects, research, policy and advocacy efforts, and future development of clinical practice guidelines that support the integration of supportive care throughout the continuum of kidney disease.

Background and Evidence

- More recently, the term “supportive care” has been used instead of “palliative care” as it is generally preferred by patients and health care professionals. Supportive care generally refers to the care that the nephrology team provides, while palliative care refers to the care provided by specialists in palliative care (Davison et al., 2024).
- Approximately 14% of people needing palliative care actually receive it (WHO, 2020).
- Older adults experience higher symptom burden and higher mortality on dialysis, and represent the growing number of dialysis patients, with median age 74 (Chou et al., 2023).
- Barriers to palliative care include cultural and social beliefs, and lack of awareness and misconceptions of what palliative care is (WHO, 2020).
- Facilitators to palliative care include specific resources, training, and support to enhance primary palliative care skills (de Barbieri et al., 2022).
- Significant policy changes are needed “to improve patient-centered care and the quality of life for seriously ill patients with kidney disease” (Diamond et al., 2021, p. 529).
- The review of the literature points to the need for healthcare providers to be competent in all aspects of comprehensive kidney supportive care, including access to quality palliative/EOL care and for all people living with CKD to have an advance medical directive. Nephrology nurses have the opportunity to improve a holistic, patient-centered approach to care by engaging in activities that support initiation of discussions around patient goals and prognosis and encouraging patient-family communication to reinforce the broader understanding of the palliative approach to ameliorate symptoms and to minimize suffering.
- ANNA supports the National Consensus Project’s *Clinical Practice Guidelines for Quality Palliative Care* (4th edition), as it relates to nephrology nursing care.

Glossary

Advance Care Planning (ACP) – an ongoing process of communication among patient, family, and friends, and the health care team in which the individual’s preferences for a surrogate and for future medical care are determined prospectively (sometimes including the completion of a written medical directive), updated periodically, and respected when the patient no longer has the capacity to participate in medical decision-making (NIA, 2022; Davison, 2026).

Advance Directive (AD)/Advance Medical Directive (AMD) – an oral or written statement by an individual with decision-making capacity expressing their preferences for a surrogate and/or future medical care in the event they become unable to participate in medical decision-making. All 50 states have one or more laws recognizing written advanced directives. There are two types of advance directives: a health care proxy and a living will. The health care proxy designates a person to make decisions if the patient loses decision-making capacity. The health care proxy is known in some states as a medical power of attorney or a durable power of attorney for health care. The living will, also known as an instruction directive, indicates the patient’s wishes that are to be followed if they lose decision-making capacity. Wishes may refer to care in the event of medical conditions such as terminal illness or a persistent vegetative state. In some states, both functions are combined in the living will (NIA, 2022).

End-of-life (EOL) care – support and medical care provided to those who are terminally ill at the time surrounding death (NIA, 2022).

Nephrology Nurse's Role in Kidney Supportive Care, including Palliative and End-of Life Care

Hospice – a team approach to treating the terminally ill individual, usually in the home, that uses the principles of palliative care to help meet the physical, psychological, social, and spiritual needs of the individual and family. Hospice treats the person not the disease; considers the entire family the unit of care; and provides bereavement counseling for the family after the individual's death (Davison, 2026).

Kidney supportive care – involves the components of shared decision-making, symptom management, crisis planning, advance care planning, spiritual care, integration with community services, and end-of-life care considerations and bereavement (Davison et al., 2024), which includes palliative care and end of life care.

Palliative care – patient- and family-centered approach to improve quality of life in persons with life-threatening illnesses that “prevents and relieves suffering through the early identification, correct assessment and treatment of pain and other problems, whether physical, psychosocial or spiritual” (WHO, 2020, para 1).

Shared decision-making – “Is an essential phase of the communication pathway and embedded in serious illness treatment processes. This interactional process provides a vehicle to honor both the provider's expertise and the patient's right to be fully informed of all care options, and their potential harms and benefits. Health care decisions are collaboratively defined through consideration of the best scientific evidence available, as well as the patient's values and preferences” (Fink-Samnack, 2026).

References

- de Barbieri, I., Strini, V., Noble, H., Amatori, S., & Sisti, D. (2022). Nurse-perceived facilitators and barriers to palliative care in patients with kidney disease: A European Delphi survey. *Journal of Renal Care*, 48(1), 49–59. <https://doi.org/10.1111/jorc.12371>
- Chou, A., Li, C., Farshid, S., Hoffman, A., & Brown, M. (2023). Survival, symptoms and hospitalization of older patients with advanced chronic kidney disease managed without dialysis. *Nephrology Dialysis Transplantation*, 38(2), 405–413. <https://doi.org/10.1093/ndt/gfac154>
- Davison, S.N. (2026) Kidney palliative care: Principles, benefits, and core components. *UpToDate*. Retrieved July 11, 2026 at <https://www.uptodate.com/contents/kidney-palliative-care-principles-benefits-and-core-components>
- Davison, S., Pommer, W., Brown, M., Douglas, C., Gelfand, S., Gueco, I., Hole, B., Homma, S., Kazancioglu, R., Kitamura, H., Koubar, S., Krause, R., Li, K., Lowney, A., Nagaraju, S., Niang, A., Obrador, G., Ohtake, Y., Schell, J., Scherer, J....Brennan, F. (2024). Conservative kidney management and kidney supportive care: Core components of integrated care for people with kidney failure. *Kidney International*, 105(1), 35–45. <https://doi.org/10.1016/j.kint.2023.10.001>
- Diamond, L., Armistead, N., Lupu, D., & Moss, A. (2021). Recommendations for public policy changes to improve supportive care for seriously ill patients with kidney disease. *American Journal of Kidney Disease*, 77(4), 529–537. <https://doi.org/10.1053/j.ajkd.2020.09.020>
- Fink-Samnack, E. (2026). Shared decision making in palliative care: How research informs practice for case management. *Case Management Society of America (CMSA) Today*. Retrieved on July 11, 2026 from [Shared Decision Making in Palliative Care: How Research Informs Practice for Case Management - Case Management Society of America \(cmsatoday.com\)](https://www.cmsatoday.com/Shared-Decision-Making-in-Palliative-Care-How-Research-Informs-Practice-for-Case-Management)

- Gelfand, S., Mandel, E., Mendu, M., & Lakin, J. (2020). Palliative care in the Advancing American Kidney Health Initiative: A call for inclusion in kidney care delivery models. *American Journal of Kidney Diseases*, 76(6), 877-882. <https://doi.org/10.1053/j.ajkd.2020.07.022>
- National Consensus Project for Quality Palliative Care. (2018) *Clinical practice guidelines for quality palliative care* (4th edition). National Coalition for Hospice and Palliative Care. Retrieved July 3, 2024 from https://www.nationalcoalitionhpc.org/wp-content/uploads/2020/07/NCHPC-NCPGuidelines_4thED_web_FINAL.pdf
- National Institute on Aging (NIA). (2022). *Advance care planning*. National Institutes of Health. <https://www.nia.nih.gov/health/advance-care-planning/advance-care-planning-advance-directives-health-care>
- World Health Organization (WHO). (2020). *Palliative care*. <https://www.who.int/news-room/fact-sheets/detail/palliative-care>

Suggested Readings

- American Nurses Association. (2020). *Position statement. Nursing care and do-not-resuscitate (DNR) decisions*. Retrieved from <https://www.nursingworld.org/~494a87/globalassets/practiceandpolicy/nursing-excellence/ana-position-statements/social-causes-and-health-care/nursing-care-and-do-not-resuscitate-dnr-decisions-final-nursingworld.pdf>
- American Nurses Association. (2025). *Code of ethics for nurse*.
- Centers for Medicare and Medicaid Services. (2018). *Hospice, end of life, and/or palliative care critical element pathway*. Retrieved July 3, 2024 from <https://www.cms.gov/files/document/cms-20073-hospice-end-lifepdf>

Suggested Tools

- Edmonton Symptom Assessment-Renal:
<https://www.albertahealthservices.ca/frm-20351.pdf>
- Integrated Palliative Care Outcome Scale-Renal (IPOS-Renal):
<https://pos-pal.org/maix/ipos-renal-in-english.php>
- Predicting 6 and 12 month mortality in CKD patients:
https://qxmd.com/calculate/calculator_446/predicting-6-and-12-month-mortality-in-ckd-patients

Adopted by the ANNA Board of Directors in August 2024.

ANNA Position Statements are reviewed and reaffirmed biennially.

Most recently revised: August 2026